

ProSci Incorporated 12170 Flint Place Poway, CA 92064, USA prosci-inc.com P: +1 (888) 513-9525 Local: +1 (858) 513-2638 Fax: +1 (858) 513-2692

sRANK Ligand Antibody


CATALOG NUMBER: 3963

Specifications

Host Species	Rabbit
Species Reactivity	Human, Mouse, Rat
Immunogen	Anti-sRANK-L antibody (3963) was raised against a peptide corresponding to 14 amino acids near the center of human sRANK-L. The immunogen is located within amino acids 140-190 of sRANK-L.
Conjugate	Unconjugated
Tested Applications	ELISA, IHC-P, WB
User Note	Optimal dilutions for each application to be determined by the researcher.
Predicted Molecular Weight	Predicted: 36kD Observed: 36 kD


Properties

Purification	sRANK Ligand Antibody is affinity chromatography purified via peptide column.
Clonality	Polyclonal
Isotype	IgG
Physical State	Liquid
Buffer	sRANK Ligand Antibody is supplied in PBS containing 0.02% sodium azide.
Concentration	1 mg/mL
Storage Conditions	sRANK Ligand antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.




sRANK Ligand Antibody 1

Figure 1 Western Blot Validation in Human Cell Lines Loading: 15 μg of lysates per lane.Antibodies: sRANK-L 3963 (1 μg/mL), 1h incubation at RT in 5% NFDM/TBST.Secondary: Goat ...




sRANK Ligand Antibody 3

Figure 3 Western Blot Validation with Recombinant Protein Loading: 30 ng of human sRANK-L recombinant protein per lane.Antibodies: sRANK-L 3963 (A: 1 μg/mL, B: 2 μg/mL and C: 4...



sRANK Ligand Antibody 2

Figure 1 Western Blot Validation in Mouse TissuesLoading: 15 μg of lysates per lane.Antibodies: sRANK-L 3963 (2 μg/mL), 1h incubation at RT in 5% NFDM/TBST.Secondary: Goat anti...



sRANK Ligand Antibody 4

Figure 4 Western Blot Validation in Rat Liver Tissue Loading: 15 μ g of lysates per lane.Antibodies: sRANK-L 3963 (A: 0.25 μ g/mL and B: 0.5 μ g/mL), 1h incubation at RT in 5% NFDM...

Disclaimer

DisclaimerOptimal dilutions/concentrations should be determined by the end user. The information provided is a guideline for product use. This product is for research use only.

For research use only. For additional information, visit ProSci's Terms and Conditions Page.